

Soins primaires, pharmacies et interventions sur le risque alcool

Philippe Michaud
Paris, 25 novembre 2014




Centre Victor-Segalen
(CSAPA de Clichy, 92)



A problème fréquent réponse nombreuse

- 5 millions de personnes qui consomment de l'alcool de façon dangereuse
- 270 CSAPA...
- 625 services d'urgences (15 millions de passages)
- 100 000 médecins généralistes
- 54 000 pharmaciens en officine
- 7 300 infirmières scolaires

Les cinq premiers facteurs de risque de morbidité et de mortalité prématurée en Europe



Source : OMS, 2002

RPIB en pharmacies

- Watson MC, Blenkisopp A. The feasibility of providing community pharmacy-based services for alcohol misuse: a literature review. *Int J Pharm Pract* 2009; 17 (4): 199-205.
- Dhital R., Whittlesea CM, Norman IJ, Milligan P. Community pharmacy service users' views and perceptions of alcohol screening and brief intervention. *Drug Alcohol Rev.* 2010; 29 (6): 596-602.
- Sheridan J, Smart R, McCormick R. Estimating problem drinking among community pharmacy customers: what did pharmacists think of the method? *Int J Pharm Practice.* 2010; 18: 290-296.
- Sheridan J, Stewart J, Smart R, McCormick R. Risky drinking among community pharmacy customers in New Zealand and their attitudes toward pharmacist screening and brief interventions. *Drug Alcohol Rev.* 2011.